

**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1999

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- Coloured covers / Couverture de couleur
- Covers damaged / Couverture endommagée
- Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- Cover title missing / Le titre de couverture manque
- Coloured maps / Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- Bound with other material / Relié avec d'autres documents
- Only edition available / Seule édition disponible
- Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- Additional comments / Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured pages / Pages de couleur
- Pages damaged / Pages endommagées
- Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- Pages detached / Pages détachées
- Showthrough / Transparence
- Quality of print varies / Qualité inégale de l'impression
- Includes supplementary material / Comprend du matériel supplémentaire
- Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below / Ce document est filmé au taux de réduction indiqué ci-dessous.

10x	14x	18x	22x	26x	30x
12x	16x	20x	24x	28x	32x

The copy filmed here has been reproduced thanks to the generosity of:



École polytechnique,
Université de Montréal,
Bibliothèque

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ▽ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

École polytechnique,
Université de Montréal,
Bibliothèque

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.


Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ▽ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

MICROCOPY RESOLUTION TEST CHART

ANSI and ISO TEST CHART No. 2



APPLIED IMAGE Inc

Applied Image Inc.
1000 Corporate Park Drive
San Jose, California 95134
(408) 966-9300



PUBLICATIONS
OF THE
DOMINION OBSERVATORY

PUBLICATIONS

OF THE

DOMINION OBSERVATORY

OTTAWA, CANADA

Vol. IV, No. 1

16262

ORBIT OF THE SPECTROSCOPIC BINARY χ AURIGAE

AND A NOTE ON 12 LACERTAE

BY REYNOLD K. YOUNG, Ph.D.

The binary character of χ Aurigae ($\alpha = 5^{\text{h}} 26^{\text{m}}$, $\delta = +32^{\circ} 7'$, mag. 4.88, type B1) was announced by Frost and Adams in the *Astrophysical Journal* in 1903. An orbit has been determined from eighty-eight single-prism spectrograms secured at this observatory during the years 1913, 1914, 1915, and 1916. An idea of the spectrum may be obtained from Table I, which gives the wave-lengths of the lines used in making the reductions, the number of times each line was measured and other data necessary for correcting the wave-lengths by the usual method of making the weighted mean of the residuals equal to zero. The residuals quoted are taken in the sense, observed minus the mean of the plate. In addition to the lines given in this table, the *H* and *K* lines of calcium are present and generally were measured, but they were not included with the rest of the lines for reasons which will appear later. On well-exposed plates all of the lines given are fairly sharp. Those due to silicon and some of those due to helium are faint, which accounts for the small number of times they were measured.

The velocities obtained at the Yerkes and Ottawa observatories follow in Tables II and III. The headings of the columns in these tables are self-explanatory. The residuals, as given under O-C, were obtained graphically from the final curve, and so may occasionally be a tenth of a kilometre or so in error.

TABLE I

Element	Wave-Length	Arithmetic Residual	Algebraic Residual	Total Weight	Number of Times Measured
Helium.....	3964.875	7.5	+2.1	40½	32
Helium.....	4009.417	6.8	+2.4	8	23
Helium.....	4026.352	5.7	+2.9	39½	79
Hydrogen.....	4101.890	5.8	-0.6	30½	64
Helium.....	4121.016	8.3	-0.9	5½	19
Silicon.....	4128.211	10.0	-3.2	9	29
Silicon.....	4131.047	6.2	-2.7	7½	24
Helium.....	4143.928	7.7	-3.1	22½	61
Carbon.....	4267.301	7.1	+2.8	13½	40
Hydrogen.....	4340.634	6.7	-3.0	48½	83
Helium.....	4388.100	7.2	-4.3	37	76
Helium.....	4471.676	9.1	+3.2	65	86
Magnesium.....	4481.400	5.7	-1.2	34	72

D

CAL

MT 403

PO4-01

54002-1

ECOLE POLYTECHIQUE BIBLIOTHEQUE

2

PUBLICATIONS OF THE DOMINION OBSERVATORY

TABLE II
YERKES OBSERVATIONS OF X AFRIGE

	Date	Julian Day	Velocity	O-C
1903				
Sept. 5		2,416,363.825	+24.2	+12.8
Oct. 10		298.782	+12.2	-2.4
Oct. 16		401.841	+13.8	-1.2

TABLE III
OTTAWA OBSERVATIONS OF X AURIGE

Plate	Observer*	Date	Julian Day	Velocity	Weight	O-C	Velocity <i>H</i> and <i>K</i> Lines	Weight
		1913						
5791	P	Nov. 4	2,420,076.915	-19.8	1	+ 0.5	-10.6	1
5798	Y	Nov. 5	077.872	-17.5	1	+ 2.7	-11.8	1
5804	Y	Nov. 6	078.719	-20.6	1	- 0.5
5818	P	Dec. 8	110.774	-19.2	1	- 3.4	-12.7	1
5834	Y	Dec. 18	120.628	- 9.8	1	+ 4.5	- 9.5	1
5849	C	Dec. 22	124.780	-22.3	1	- 8.7	- 1.9	1
		1914						
5873	Y	Jan. 1	134.729	-14.6	1	- 2.7
5894	P	Jan. 21	151.705	-14.5	1	- 5.9	+ 2.7	1
5922	P	Feb. 9	173.689	-15.2	2	- 9.7
5925	Y	Feb. 11	175.663	+ 1.5	1	+ 6.7
5939	Y	Feb. 15	179.589	- 2.5	1	+ 2.0
5957	C-P	Feb. 23	187.665	+ 6.7	1	+ 9.9	+ 1.8	1
5973	Y	Mar. 11	203.576	- 8.4	1	- 7.7	+ 1.3	1
6397	Y	Sept. 17	393.889	+ 7.6	1	-10.2	- 6.0	1
6460	Y	Oct. 1	407.839	+17.6	1	- 0.3	+ 8.9	1
6470	C	Oct. 2	408.833	+29.1	1	+11.3	+15.2	1
6483	H	Oct. 4	410.822	+18.0	1	+ 0.4	+12.5	1
6486	Y	Oct. 6	412.914	+22.2	1	+ 4.3	+16.9	1
6489	Y	Oct. 11	417.887	+21.1	1	+ 3.3	+11.0	1
6508	H	Oct. 20	426.791	+19.0	1	+ 1.3
6526	Y	Oct. 22	428.914	+24.7	1	+ 7.1	+13.6	1
6562	H	Nov. 17	451.824	+17.8	1	+ 1.2	+ 9.9	1
6587	P-Y	Nov. 28	465.748	+ 5.2	1	-12.5	+ 3.2	1
6593	C-P	Dec. 4	471.757	+20.8	1	+ 5.8	+12.3	1
6599	Y	Dec. 5	472.759	+16.4	1	+ 1.4	+17.5	1
6604	Y-H	Dec. 6	473.627	+13.3	1	- 1.6	+ 7.1	1
6617	C-P	Dec. 14	478.733	+10.7	1	- 3.7	+10.5	1
6627	Y	Dec. 15	482.637	+14.6	1	+ 0.6	- 3.7	1
6639	P	Dec. 16	483.757	+15.8	1	+ 2.0	- 0.1	1

ORBIT OF THE SPECTROSCOPIC BINARY X AURIGE

TABLE III
OTTAWA OBSERVATIONS OF α AURIGE—Continued

Plate	Observer*	Date	Julian Day	Velocity	Weight	O. C.	Velocity <i>H</i> and <i>K</i> Lines	Weight
1914								
6643	H	Dec. 17	2,420,484.654	+14.5	1	+ 0.8	+11.2	1
6646	Y	Dec. 20	2,420,496	+15.3	1	+ 1.9	+3.4	1
6652	Y	Dec. 22	2,420,502	+10.1	1	- 2.8	+6.2	1
6667	Y	Dec. 30	2,420,571	+16.2	1	+ 3.9	+8.9	1
1915								
6680	C	Jan. 4	502,714	+11.7	1	+ 0.2	+6.9	1
6702	Y	Jan. 10	508,592	+ 1.1	1	- 9.1	+7.2	1
6731	Y	Jan. 24	522,609	+ 7.5	1	- 0.5	+3.6	1
6750	H-Y	Jan. 28	526,721	+ 8.1	1	+ 1.0	+6.7	1
6783	C	Feb. 12	541,610	+ 6.2	1	+ 2.1	+9.3	1
6788	H	Feb. 17	546,599	+11.2	1	+ 8.3	+8.8	1
6819	Y	Feb. 28	557,604	- 2.8	1	- 3.2	-1.1	1
6841	Y	Mar. 7	564,573	+ 1.7	1	+ 3.2	-2.6	1
6879	Y	Mar. 19	576,555	-11.3	1	- 6.8	+4.8	1
6896	Y	Mar. 30	587,528	- 4.8	1	+ 2.3	0.0	1
6919	Y	Apr. 13	601,536	- 6.8	1	+ 1.1	-5.6	1
6956	Y	Apr. 28	616,556	-17.3	1	- 2.7	-18.0	1
7135	Y	July 29	707,858	-21.8	1	+ 0.7	1	1
7152	H	Aug. 10	720,851	-24.5	1	- 3.2	1	1
7175	Y	Aug. 26	736,826	-18.7	1	+ 0.9	1	1
7199	Y	Sept. 2	743,883	-21.6	1	- 2.8
7227	Y	Sept. 9	750,906	-16.8	1	+ 1.1
7264	Pm	Sept. 17	758,875	-11.8	1	+ 5.0
7280	H	Sept. 21	762,900	-13.9	1	+ 2.2	-3.7	1
7316	Y	Sept. 30	771,881	-22.2	1	- 7.1	-11.7	1
7323	Y	Oct. 9	780,816	- 7.1	1	+ 6.3	-12.9	1
7345	C	Oct. 15	786,816	- 2.2	1	+10.3
7364	Y	Oct. 24	795,892	-12.2	1	- 1.2	-0.6	1
7373	Y	Nov. 3	805,622	- 9.8	1	- 0.3	-6.6	1
7384	Y	Nov. 6	808,862	-11.6	1	- 5.7
7411	H	Nov. 12	814,740	- 5.3	1	+ 2.6	+10.1	1
7415	C	Nov. 17	819,625	-16.6	1	- 9.5
7429	Y-C	Nov. 24	826,708	- 1.4	1	+ 4.4	-2.4	1
7440	Y	Dec. 3	835,570	+ 6.4	1	+10.7
7447	C	Dec. 10	842,622	- 4.9	1	- 1.7	+1.5	1
7454	P	Dec. 20	852,649	- 9.2	1	- 7.7	-3.9	1
7458	Y	Dec. 28	860,651	+ 3.0	1	+ 3.2	-8.0	1
1916								
7475	C	Jan. 7	870,514	- 5.8	1	- 7.0	+4.5	1
7481	H	Jan. 13	876,569	- 2.0	1	- 4.0	+1.8	1
7491	Y	Jan. 28	891,574	+ 2.6	1	- 1.6	+0.3	1
7502	H	Feb. 3	897,728	+12.6	1	+ 7.6	+12.6	1
7504	Y	Feb. 10	904,584	+ 6.2	1	+ 0.2	+11.6	1
7510	P	Feb. 19	913,669	+13.5	1	+ 6.3
7512	Y	Feb. 20	914,501	+ 9.4	1	+ 3.1	+5.3	1
7517	C	Feb. 23	917,629	+ 6.5	1	- 1.1	+8.5	1
7527	Y	Feb. 29	923,524	+ 4.8	1	- 6.6	+5.4	1

PUBLICATIONS OF THE DOMINION OBSERVATORY

TABLE III
OTTAWA OBSERVATIONS OF α AURIGE—*Concluded*

Plate	Observer*	Date	Julian Day	Velocity	Weight	O-C	Velocity <i>H</i> and <i>K</i> Lines	Weight
		1916						
7532	C	Mar. 1	2,420,921.501	+ 4.8	1	- 3.7	- 2.0	1
7536	H	Mar. 2	925.621	+ 8.5	1	- 0.1	+ 3.9	1
7539	Y	Mar. 5	928.531	+ 0.9	1	- 8.1	+ 6.1	1
7542	C	Mar. 10	933.535	+10.8	1	- 1.3	+10.1	1
7549	Y	Mar. 17	940.597	+11.5	1	+ 1.2	+ 4.2	1
7550	Y	Mar. 17	940.642	+ 8.8	1	- 1.5	+ 5.5	1
7558	Y-H	Mar. 19	942.637	+ 8.5	1	- 2.1	+ 9.2	1
7562	Y	Mar. 21	944.510	+14.0	1	+ 3.2	+11.7	1
7571	H	Mar. 23	946.618	+14.8	1	+ 3.8	+ 6.5	1
7585	Y	Mar. 30	953.575	+ 7.8	1	- 4.0	+ 7.8	1
7591	Y	Apr. 2	956.572	+ 9.0	1	- 3.0	+ 3.7	1
7601	H	Apr. 5	959.525	+16.5	1	+ 4.2	+13.0	1
7624	Y	May 2	986.548	+ 9.3	1	- 5.5	+ 0.3	1
7632	H	May 4	988.555	+26.1	1	+11.2	+17.4	1

*C=Cannon; H=Harper; P=Parker; P=Plaskett, J. S.; P^{II}=Plaskett, H. H.; Y=Young

ORBIT OF THE SPECTROSCOPIC BINARY X AURIGE

MEASURES OF X AURIGE

X	5791		5798		5804		5818		5834		5849		5873	
	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.
3933.825	-29.1	1	-29.9	1	-	-	-15.4	1	-7.2	1	+2.8	1	-	-
3964.875	-31.7	1	-	-	-	-	-	-	-8.0	1	-	-	-	-
4009.417	-49.0	1	-	-	-	-	-22.9	1	-3.1	1	-15.3	1	+3.8	1
4026.352	-33.1	1	-18.0	1	-35.1	1	-26.0	1	-22.5	1	-	-	-	-
4101.890	-35.6	1	-40.4	1	-26.0	1	-25.4	1	-	-	-	-	-	-
4121.016	-25.4	1	-15.0	1	-15.9	1	-	-	-	-	-	-	-	-
4128.214	-53.0	1	-	-	-	-	-	-	-	-	-	-	-	-
4131.047	-31.3	1	-	-	-	-	-36.7	1	-10.8	1	-	-	-	-
4143.928	-10.0	1	-24.8	1	-	-	-	-	-	-	-	-	-	-
4267.301	-	-	-28.5	1	-48.2	1	-	-	-18.8	1	-9.1	1	-15.0	1
4340.634	-35.8	1	-33.5	1	-	-	-	-	-	-	-43.5	1	+10.5	1
4388.100	-38.2	1	-10.6	1	-17.8	1	-	-	-	-	-22.3	1	-2.5	1
4471.676	-53.1	1	-31.6	1	-41.3	1	-31.5	1	-	-	-	-	-15.1	1
4481.400	-31.9	1	-42.3	1	-41.6	1	-	-	-	-	-	-	-	-
Weighted mean	-38.40		-35.60		-38.48		-21.88		-7.47		-17.68		-2.42	
V _a	+18.94		+18.57		+18.21		+3.07		-2.06		-4.22		-9.30	
V _d	-0.41		-0.15		-0.07		-0.07		0.00		-0.46		-0.42	
Curv.	-0.28		-0.28		-0.28		-0.28		-0.28		-0.28		-0.28	
Radial Velocity <i>H</i> and <i>K</i>	-19.8		-17.5		-20.6		-19.2		-9.8		-22.3		-12.1	
	-10.6		-11.8		-		-12.7		-9.5		-1.9		-	

MEASURES OF X AURIGE.—Continued

λ	5891		5922		5925		5939		5957		5973		6397	
	Vel.	Wt.												
3933.825	+21.4	1							+30.1	1	+31.3	1	-35.0	1
3961.875							+21.2	1					-25.5	1
3968.625													+6.9	1
4026.352	+11.1	1			+27.2	1	+32.5	1	+28.1	1	+22.8	1	+5.6	1
4101.890	+1.7	1			+37.6	1	+16.0	1			+22.6	1		
4121.016					+42.4	1	+5.7	1						
4128.211							+4.8	1						
4131.047							+22.1	1						
4143.928							+31.1	1	+22.4	1	+13.6	1		
4267.301							+30.1	1	+28.9	1	+21.5	1	-33.0	1
4340.634	-7.9	1	+11.1	1	+43.4	1	+37.6	1	+36.1	1	+19.3	1	-27.0	1
4388.400	+22.1	1	+24.7	1	+17.7	1	+27.1	1	+27.1	1	+16.5	1	-20.1	1
4471.676	+12.5	1	+2.5	1	+21.2	1	+13.7	1	+44.5	1	+33.7	1	-22.3	1
4481.400	-5.0	1			+47.6	1	+10.1	1	+41.1	1	+15.2	1	-21.9	1
Weighted mean	+4.18		+10.26		+27.52		+24.48		+35.30		+21.55		-21.46	
V_a	-18.42		-25.00		-25.54		-26.60		-28.08		-29.54		+29.22	
V_d	0.00		-0.20		-0.18		-0.10		-0.22		-0.15		+0.11	
Curv	-0.28		-0.28		-0.28		-0.28		-0.28		-0.28		-0.28	
Radial Velocity	-11.5		-15.2		+1.5		-2.5		+6.7		-8.1		+7.6	
H and K	+2.7								+1.8		+1.3		-6.0	

MEASURES OF X-AFRIGA. *Continued*

λ	6460		6470		6483		6486		6489		6508		6526	
	Vel	Wt												
3931.825	-19.4	4	-12.7	4	-17.5	4	-8.7	4	-15.4	4	-22.9	4	-9.5	4
3964.875					-14.0	4	-9.8	4			+3.4	4	-20.5	4
3998.625									0.0	4	-16.5	4	+1.7	4
4009.447	-9.6	4	+10.2	4	-22.7	4	+3.4	4			-17.6	4	+3.8	4
4026.352	0.0	4	+0.9	4	-16.6	4	-22.3	4			+4.7	4	-9.4	4
4101.890	-6.5	4	-6.5	4									+12.3	4
4121.016			-4.7	4					-1.8	4				
4128.244	-22.8	4												
4131.047	-18.4	4							-11.5	4			+9.6	4
4143.928	-5.9	4	+5.8	4	-10.5	4	-1.9	4					+7.4	4
4267.301					-19.1	4	-16.9	4	-10.1	4	-3.8	4	-9.0	4
4340.634	-21.4	4	+5.6	4							+12.8	4	-7.0	4
4388.100	-8.2	4	0.0	4							+1.2	4	+7.4	4
4471.676	-2.5	4	-1.2	4	+3.7	4	+6.2	4	-3.7	4				
4481.400	-7.4	4	-2.5	4	-5.0	4	-9.9	4	-1.2	4				
Weighted mean	-10.53		+1.24		-9.64		-3.82		-5.08		-5.01		+1.59	
V_x	+28.48		+28.05		+27.75		+26.30		+26.43		+24.19		+23.50	
V_d	+0.42		+0.14		+0.14		-0.07		0.00		+0.11		-0.11	
Curv.	-0.28		-0.28		-0.28		-0.28		-0.28		-0.28		-0.28	
Radial Velocity H and K	+17.6		+29.1		+18.0		+22.2		+21.1		+19.0		+24.7	
	+8.9		+15.2		+12.5		+16.9		+11.0				+13.6	

MEASURES OF α AFRICAE - *Continued*

λ	6562		6587		6593		6599		6604		6617		6627	
	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.
3933+825	+ 2.4	1	+ 4.8	1	+11.2	1	+ 8.0	1	+ 3.2	1	+ 8.8	1	- 7.2	1
3964+875	+18.1	1	+11.0	1	-	-	+23.9	1	+ 8.2	1	+ 5.8	1	-	-
3968-625	+ 5.0	1	-	-	+ 3.5	1	+18.2	1	-	-	+ 9.9	1	+ 5.8	1
4009-417	-	-	+18.9	1	+12.9	1	-	-	+18.9	1	-	-	-	-
4026+352	+ 3.5	1	+ 0.8	1	+11.3	1	+16.2	1	+18.3	1	+22.6	1	+13.1	1
4101+890	-	-	+ 6.5	1	+21.4	1	- 0.0	1	+ 1.9	1	+ 6.5	1	+31.7	1
4121+016	-	-	-	-	+ 6.6	1	-	-	-	-	-	-	-	-
4128+211	-	-	+ 1.8	1	+23.8	1	-	-	-	-	-	-	-	-
4131+017	-	-	+ 1.9	1	-	-	+15.3	1	-	-	-	-	-	-
4143+928	- 1.8	1	-16.4	1	+28.0	1	- 1.9	1	- 2.9	1	+ 1.0	1	+ 1.0	1
4267-301	-	-	-	-	+22.4	1	-	-	+ 6.1	1	-	-	+31.0	1
4340-631	+20.3	1	-13.6	1	+18.1	1	- 4.5	1	+ 4.5	1	+10.2	1	+ 6.8	1
4388-100	-	-	-	-	+ 1.7	1	+35.1	1	+11.7	1	+19.9	1	+10.5	1
4471+676	+ 6.2	1	+24.9	1	- 9.9	1	+33.6	1	+ 6.8	1	+ 8.7	1	+11.2	1
4481+400	+16.2	1	+11.3	1	+21.3	1	+22.5	1	+ 8.7	1	+ 6.3	1	+11.0	1
Weighted mean	+ 9.39	- 2.88	+ 15.85	- 11.91	+ 9.41	- 9.27	+ 15.43	-	-	-	-	-	-	-
V_a	+ 8.83	+ 8.32	+ 5.28	+ 1.77	+ 4.32	+ 1.78	- 0.42	-	-	-	-	-	-	-
V_d	- 0.07	- 0.00	- 0.04	- 0.03	- 0.16	- 0.04	+ 0.12	-	-	-	-	-	-	-
Cury	- 0.28	- 0.28	- 0.28	- 0.28	- 0.28	- 0.28	- 0.28	-	-	-	-	-	-	-
Radial Velocity	+ 17.8	+ 5.2	+ 20.8	+ 16.4	+ 13.3	+ 10.7	+ 14.6	-	-	-	-	-	-	-
H and K	+ 9.9	+ 3.2	+ 12.3	+ 17.5	+ 7.1	+ 10.5	+ 3.7	-	-	-	-	-	-	-

MEASURES OF X AURIGÆ—Continued

λ	6639		6643		6646		6652		6667		6680		6702	
	Vel.	Wt.												
3933.825	+ 0.8	1/2	+15.2	1/2	+ 6.4	1/2	+10.4	1	+16.8	1	+ 9.6	1	+23.2	1
3964.875	+17.3	1/2	+20.6	1/2	+15.6	1/2	+15.6	1/4
3968.625	+ 3.3	1/2	+10.7	1/2	+10.1	1/2	+17.1	1	+27.2	1/4	+16.5	1/4
4009.417	+ 9.4	1/4	+17.2	1/4	+28.3	1/4	+25.3	1
4026.352	+17.4	1/2	+25.3	1/2	+18.3	1/2	+16.6	1/2	+35.7	1/2	+ 2.8	1
4101.890	+23.4	1/4	+14.9	1/2	+28.0	1/4	+21.4	1/4
4121.016	+17.1	1/4	+18.9	1/4
4128.211	0.0	1/4	+ 7.6	1/4	+ 3.8	1	+22.9	1
4131.047	+10.5	1/4	+ 7.6	1/4	+15.3	1/4	+26.8	1
4143.928	+13.5	1/2	+22.2	1/4	+21.3	1	+ 5.8	1/4
4267.301	+17.1	1/4	+20.3	1/4	+16.0	1	+13.9	1/2
4340.634	+26.0	1/2	+ 7.9	1/2	+15.9	1/2	+20.3	1/2	+17.0	1	+14.7	1/4	+ 5.7	1
4388.100	+ 9.4	1/4	+28.0	1/4	+28.1	1/4	+ 1.2	1/2	+26.3	1/2	+18.1	1/4	+ 9.4	1/4
4471.676	+18.7	1	+14.9	1/4	+ 6.7	1/2	+18.0	1/2	+31.7	1	+33.6	1/4	+18.7	1
4481.400	+ 2.6	1/4	+13.8	1	+31.3	1/2	+22.5	1/2	+23.8	1/2	+28.2	1/4	+17.5	1/2
Weighted mean	+ 17.23	+ 16.16	+ 18.29	+ 14.47	+ 21.41	+ 23.19	+ 15.18
V_a	- 0.96	- 1.42	- 2.91	- 3.99	- 8.09	- 10.62	- 13.56
V_d	- 0.15	0.00	+ 0.21	+ 0.14	+ 0.15	- 0.15	+ 0.07
Curv.	- 0.28	- 0.28	- 0.28	- 0.28	- 0.28	- 0.28	- 0.28
Radial Velocity H and K	+ 15.8	+ 14.5	+ 15.3	+ 10.4	+ 16.2	+ 11.7	+ 1.4
	- 0.1	+ 11.2	+ 3.1	+ 6.2	+ 8.9	+ 6.9	+ 7.2

MEASURES OF X AURIGAE - *Continued*

λ	6734		6750		6783		6788		6819		6841		6879	
	Vel.	Wt.												
3933-825	+22.4	1/2	+27.2	1	+32.0	1	+36.0	1	+30.4	1	+27.2	1	+30.4	1/2
3964-875	+21.4	1	+21.4	1	+33.7	1	+33.7	1	+30.0	1	+38.8	1	+38.8	1
3968-625	+29.7	1/2	+42.1	1/2	+36.3	1/2	+19.0	1	+41.2	1	+31.0	1/2	+24.4	1/2
4009-417	+21.9	1/2	+37.8	1	+27.0	1	+41.9	1/2	+26.2	1	+21.4	1/2	+6.5	1
4026-352	+11.4	1	+42.7	1	+9.3	1	+11.0	1	+29.9	1/2	+43.9	1/2	+21.4	1
4101-890	+37.0	1	+35.1	1	+9.3	1	+11.0	1	+21.4	1/2	+43.8	1	+21.4	1
4121-016	+37.0	1	+35.1	1	+9.3	1	+11.0	1	+21.4	1/2	+43.8	1	+21.4	1
4128-211	+25.8	1	+21.3	1	+18.3	1	+13.5	1	+26.7	1	+43.8	1	+21.4	1
4131-047	+19.3	1/2	+21.3	1	+18.3	1	+13.5	1	+21.5	1	+22.6	1	+20.4	1
4143-928	+24.9	1/2	+23.8	1	+30.5	1	+30.5	1	+21.5	1	+28.1	1	+7.0	1
4267-301	+24.6	1	+35.1	1/2	+31.7	1/2	+35.1	1	+12.7	1	+34.8	1	+18.0	1
4340-634	+23.8	1	+36.1	1	+31.1	1	+42.3	1	+31.1	1	+29.4	1	+29.4	1
4388-100	+21.1	1/2	+36.1	1	+31.1	1	+42.3	1	+31.1	1	+29.4	1	+29.4	1
4471-076	+37.3	1/2	+36.1	1	+31.1	1	+36.9	1	+13.1	1	+29.4	1	+27.5	1
4481-400	+23.2	1	+27.5	1/2	+11.1	1/2	+36.9	1	+13.1	1	+29.4	1	+27.5	1
Weighted mean	+ 27.34		+ 29.50		+ 32.34		+ 38.49		+ 26.34		+ 31.46		+ 18.49	
V_a	- 19.49		- 20.90		- 25.69		- 26.90		- 28.75		- 29.37		- 29.42	
V_d	- 0.02		- 0.20		- 0.16		- 0.11		- 0.16		- 0.12		- 0.16	
Curv.	- 0.28		- 0.28		- 0.28		- 0.28		- 0.28		- 0.28		- 0.28	
Radial Velocity H and K	+ 7.5		+ 8.1		+ 6.2		+ 11.2		- 2.8		+ 1.7		- 11.3	
	+ 3.6		+ 6.7		+ 9.3		+ 8.8		- 1.1		- 2.6		+ 4.8	

MEASURES OF X AURIGE—Continued

λ	6896		6919		6956		7135		7152		7175		7199	
	Vel.	Wt.												
3933.825	+28.0	1	+18.4	1	+ 3.6	1/2
3964.875	+13.2	1	+ 9.9	1
3968.625	+30.5	1/2	+28.9	1
4009.417	+18.9	1
4026.352	+34.0	1	+18.3	1/2	+19.9	1/4	-38.1	1	-52.0	1
4101.890	+21.5	1/2	+17.7	1/2	+ 6.5	1/4
4131.047	+21.0	1
4143.928	+27.0	1	+32.4	1/4	-66.4	1	-45.7	1
4267.301	- 6.4	1/4	-40.4	1
4340.634	+19.0	1/2	+11.1	1/2	+11.3	1/2	-11.6	1/4	-63.0	1/2	-50.0	1
4388.100	+22.2	1/2	- 2.3	1/2	-11.3	1/2	-45.5	1/2
4471.676	+33.6	1	+19.3	1	0.0	1/2	-45.8	1/2	-47.0	1	-27.2	1/2	-11.5	1/2
4481.400	+17.5	1/2	+20.0	1	+ 5.0	1/4	-33.7	1/4	-54.8	1/2	-45.5	1	-58.6	1/2
Weighted mean	+ 21.05		+ 18.38		+ 4.28		- 41.72		- 18.27		- 46.12		+ 50.00	
V _a	- 28.37		- 25.59		- 21.04		+ 19.95		+ 23.82		+ 27.48		+ 28.47	
V _d	- 0.16		- 0.22		- 0.28		+ 0.27		+ 0.25		+ 0.23		+ 0.16	
Curv.	- 0.28		- 0.28		- 0.28		- 0.28		- 0.28		- 0.28		- 0.28	
Radial Velocity <i>H</i> and <i>K</i>	- 4.8		- 6.8		- 17.3		- 21.8		- 21.5		- 18.7		- 21.6	
	0.0		- 5.6		- 18.0		

MEASURES OF χ AURIGE—Continued

λ	7227		7264		7280		7316		7323		7345		7361	
	Vel.	Wt.												
3933·825	-32·6	1	-39·8	1	-28·0	1	-23·1	1
3961·875	-28·6	1
4009·417	-36·7	1
4026·352	-35·5	1	-32·9	1	-40·7	1	-52·0	1	-30·1	1	-25·2	1	-37·3	1
4101·890	-27·9	1	-55·7	1	-32·5	1	-38·0	1
4121·016	-42·4	1
4128·211	-42·7	1	-21·3	1
4131·047	-28·5	1	-36·1	1
4113·928	-53·9	1	-48·1	1	-52·9	1	-37·7	1	-29·8	1	-38·5	1
4267·301	-50·0	1	-20·2	1	-33·9	1
4340·631	-45·0	1	-25·9	1	-50·7	1	-33·9	1	-20·3	1	-30·4	1
4388·400	-41·3	1	-47·8	1	-56·0	1	-42·0	1	-35·1	1	-29·1	1	-40·8	1
4171·676	-46·4	1	-37·7	1	-35·2	1	-48·9	1	-30·5	1	-20·3	1	-31·5	1
4181·400	-61·1	1	-61·1	1	-61·1	1	-30·6	1
Weighted mean	-45·72		-40·85		-42·79		-50·35		-33·82		-27·58		-34·68	
V_a	+ 29·06		+ 29·23		+ 29·11		+ 28·34		+ 26·90		+ 25·57		+ 23·00	
V_d	+ 0·10		+ 0·13		+ 0·01		+ 0·05		+ 0·06		+ 0·08		- 0·07	
Curv.	- 0·28		- 0·28		- 0·28		- 0·28		- 0·28		- 0·28		- 0·28	
Radial Velocity H and K	- 16·8		- 41·8		- 43·9		- 22·2		- 7·1		- 2·2		- 12·2	
	- 3·7		- 11·7		- 12·9		- 0·6	

MEASURES OF X AURIGAE -Continued

λ	7373		7384		7401		7415		7429		7440		7447	
	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.								
3933.825	-25.2	1	-0.6	1	-12.8	1	-8.0	1
3964.875	-8.2	1	-10.7	1	-17.3	1	+6.6	1
3968.625	-36.6	1	-19.1	1	-17.8	1	-1.3	1
4026.352	-29.4	1	-12.1	1	-5.1	1	-17.7	1
4101.890	-32.4	1	-26.1	1
4143.928	-45.0	1	-13.9	1
4267.301	-53.0	1	-30.0	1	-31.6	1	-3.4	1	+11.3	1	-7.9	1
4340.634	-42.9	1	-25.8	1	-31.5	1	-31.2	1	-28.1	1	0.0	1
4388.100	-37.0	1	-29.2	1	-21.8	1	-32.3	1	-6.8	1	-7.4	1	-8.7	1
4471.676	-19.9	1	-35.0	1	-3.7	1	-28.2	1	-12.5	1
4481.400	-31.3	1
Weighted mean	-28.40	1	-32.75	1	-21.10	1	-30.32	1	-11.81	1	-0.94	1	-7.11	1
V _a	+19.62	1	+18.53	1	+15.92	1	+13.76	1	+10.42	1	+7.41	1	+2.28	1
V _d	+0.22	1	-0.08	1	+0.10	1	+0.22	1	+0.05	1	+0.23	1	+0.04	1
Curv.	-0.28	1	-0.28	1	-0.28	1	-0.28	1	-0.28	1	-0.28	1	-0.28	1
Radial Velocity	-9.8	1	-14.6	1	-5.3	1	-16.6	1	-1.4	1	+6.4	1	-4.9	1
H and K	-6.6	1	+10.1	1	-2.4	1	+1.5	1

MEASURES OF X AURIGE—Continued

λ	7451		7458		7475		7481		7494		7502		7504	
	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.
3933.825	+ 0.8	1	+ 0.8	1	+ 12.8	1	+ 16.8	1	+ 20.8	1	+ 36.0	1	+ 29.6	1
3961.875	+ 7.1	1	+ 15.6	1					+ 19.8	1	+ 35.5	1	+ 46.1	1
3968.625					+ 17.1	1							+ 44.2	1
4009.417					+ 17.0	1								
4026.352	+ 3.5	1	+ 17.0	1	+ 0.1	1	+ 17.0	1	+ 25.2	1	+ 29.2	1	+ 24.0	1
4101.890	+ 15.8	1	+ 8.4	1	+ 8.1	1	+ 11.9	1	+ 25.2	1	+ 37.3	1	+ 15.9	1
4121.016									+ 30.8	1				
4128.211			+ 15.2	1			+ 23.5	1	+ 10.5	1	+ 18.1	1	+ 15.3	1
4131.017			+ 1.9	1						+ 11.0	1		+ 26.8	1
4143.928					+ 12.6	1	+ 16.4	1	+ 19.3	1			+ 30.5	1
4267.301	+ 5.3	1	+ 9.6	1	+ 24.6	1	+ 28.9	1	+ 21.6	1			+ 40.6	1
4340.634	+ 16.4	1	+ 3.4	1	+ 4.5	1	+ 13.1	1	+ 19.8	1	+ 39.0	1	+ 39.0	1
4388.400	+ 3.5	1	+ 14.0	1	0.0	1	+ 3.5	1	+ 28.1	1	+ 38.6	1	+ 38.6	1
4471.676	+ 4.3	1	+ 16.8	1	+ 3.7	1	+ 7.4	1	+ 11.3	1	+ 16.0	1	+ 32.3	1
4481.400			+ 11.3	1	+ 2.5	1	+ 8.8	1	+ 25.0	1	+ 38.8	1	+ 35.1	1
Weighted														
mean	+ 6.45		+ 10.15		+ 6.01		+ 12.98		+ 22.62		+ 35.87		+ 31.52	
V_a	+ 2.85		+ 6.97		+ 11.76		+ 11.71		+ 20.81		+ 22.72		+ 24.94	
V_d	+ 0.97		+ 0.01		+ 0.49		0.00		+ 0.02		+ 0.23		+ 0.06	
Curv.	+ 0.28		+ 0.28		0.28		+ 0.28		+ 0.28		+ 0.28		+ 0.28	
Radial														
Velocity	+ 9.2		+ 3.0		+ 5.8		+ 2.0		+ 2.6		+ 12.6		+ 6.2	
H and K	+ 3.9		+ 8.0		+ 1.5		+ 1.8		0.3		+ 12.6		+ 11.6	

MEASURES OF X AURIGE—Continued

λ	7510		7512		7517		7527		7532		7536		7539	
	Vel.	Wt.												
3933.825	+30.4	1	+34.4	4	+32.0	1	+31.2	1	+27.2	1	+28.8	1	+32.8	1
3964.875	+31.3	1	+41.6	1	+41.3	1	+38.0	1	+38.8	1
3968.625	+43.0	1	+44.2	1	+38.2	1
4009.417
4026.352	+42.3	1	+30.0	1	+35.3	1	+30.4	1	+45.3	1	+42.7	1	+31.4	1
4101.890	+45.7	1	+42.9	1	+33.6	1	+37.3	1
4128.211	+25.8	1	+33.4	1
4431.047	+19.1	1	+24.9	1
4143.928	+29.5	1	+31.9	1	+30.9	2	+19.8	1	+26.1	1
4267.301	+37.4	1	+21.6	1	+47.0	1	+24.9	1
4340.634	+41.3	1	+25.4	1	+21.5	1	+22.6	1	+25.4	1
4388.100	+32.8	1	+31.6	1	+25.8	1	+22.2	1	+28.1	1	+24.6	1
4471.676	+47.9	1	+42.9	1	+44.8	1	+36.9	1	+37.3	1	+51.6	1	+46.6	1
4481.400	+47.6	1	+30.7	1	+31.3	1	+32.6	1
Weighted mean	+ 41.20		+ 36.99		+ 34.76		+ 31.00		+ 31.60		+ 38.02		+ 30.57	
V_a	+ 27.26		+ 27.44		+ 27.98		+ 28.82		+ 28.94		+ 29.04		+ 29.29	
V_d	- 0.22		+ 0.05		- 0.05		- 0.06		0.00		- 0.19		- 0.08	
Curv.	- 0.28		- 0.28		- 0.28		- 0.28		- 0.28		- 0.28		- 0.28	
Radial Velocity H and K	+ 13.5		+ 9.4		+ 6.5		+ 1.8		+ 4.8		+ 8.5		+ 0.9	
	+ 5.3		+ 8.5		+ 5.4		- 2.0		+ 3.9		+ 6.1	

MEASURES OF X AURIGE—Continued

λ	7542		7549		7550		7558		7562		7574		7585	
	Vel.	Wt.												
3933-825	+40·0	1	+34·4	2	+36·0	1	+39·2	1	+40·0	1	+32·0	1	+37·6	1
3964-875	.	.	+42·8	1	+42·8	1	.	.	+49·3	1	+49·3	1	+39·5	1
3968-625	.	.	+33·9	2	+35·1	1	.	.	+42·1	1	+40·2	1	+35·6	1
4009-417	.	.	+28·3	1	+49·7	1
4026-352	+51·9	1	+34·0	2	+44·4	1	+34·4	1	+38·8	1	+51·5	1	+39·6	1
4101-890	+36·1	1	+43·9	2	+48·0	1	+35·5	1	+39·2	1	+44·3	1	+37·3	1
4121-016	+38·4	1
4128-214	.	.	+42·0	1	+44·8	1	+52·5	1
4131-047	.	.	+47·8	2	+31·5	1	.	.	+38·7	1
4143-928	.	.	+39·2	2	+32·9	1	+48·8	1	+18·8	1	+35·3	1	+30·4	1
4267-301	.	.	+51·3	1	+55·6	1	+52·4	1	+37·4	1
4340-634	+44·1	1	+34·5	2	+35·1	1	+30·0	1	+36·1	1	+40·7	1	+28·3	1
4388-100	+25·7	1	+42·7	2	+23·4	1	+25·8	1	+21·8	1	+39·2	1	+47·6	1
4471-676	+47·3	1	+48·5	1	+37·4	1	+52·2	1	+58·5	1	+45·4	1	+47·3	1
4481-400	.	.	+36·3	1	+35·7	1	+29·4	1	+42·6	1	+43·2	1	+42·6	1
Weighted mean	+ 40·69		+ 44·45		+ 38·84		+ 38·53		+ 43·73		+ 44·44		+ 36·52	
V_a	- 29·52		- 29·47		- 29·47		- 29·46		- 29·28		- 29·11		- 28·26	
V_d	- 0·41		- 0·22		- 0·25		- 0·25		- 0·43		- 0·24		- 0·18	
Curv.	- 0·28		- 0·28		- 0·28		- 0·28		- 0·28		- 0·28		- 0·28	
Radial Velocity	+ 10·8		+ 11·5		+ 8·8		+ 8·5		+ 14·0		+ 14·8		+ 7·8	
H and K	+ 10·1		+ 1·2		+ 5·5		+ 9·2		+ 11·7		+ 6·5		+ 7·8	

MEASURES OF X AURIGE—*Concluded*

λ	7594		7601		7624		7632					
	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.	Vel.	Wt.
3933.825	+10.0	1	+38.4	1	+20.0	1	+32.8	1
3964.875	+16.9	1	+48.6	1	+10.0	1
3968.625	+24.0	1	+45.1	1
4009.417	+33.5	1
4026.352	+31.0	1	+55.8	1	+32.7	1	+50.1	1
4101.890	+28.0	1	+55.0	1	+28.0	1	+46.6	1
4121.016	+36.5	1	+23.5	1	+52.6	1
4128.211	+24.8	1	+31.5	1	+32.5	1	+38.2	1
4131.047	+35.4	1	+36.3	1	+36.3	1
4143.928	+34.8	1	+15.0	1	+32.1	1	+49.8	1
4267.304	+56.7	1	+26.2	1
4340.634	+29.4	1	+55.4	1	+32.8	1	+28.8	1
4388.400	+51.5	1	+45.6	1	+49.2	1
4471.676	+38.6	1	+36.1	1	+51.7	1
4481.400	+42.6	1	+30.0	1	+35.1	1	+45.1	1
Weighted mean	+ 37.27		+ 44.46		+ 28.97		+ 45.05	
V_d	- 27.80		- 27.22		- 19.15		- 18.40	
V_d	- 0.23		- 0.49		- 0.28		- 0.28	
Curv.	- 0.28		- 0.28		- 0.28		- 0.28	
Radial Velocity	+ 9.0		+ 16.5		+ 9.3		+ 26.1	
H and K	+ 3.7		+ 13.0		+ 0.3		+ 17.4	

The period is remarkably long for an early type star, and it so happens that the observations taken at the Yerkes Observatory fall almost at the top of the curve, so near this point, in fact, that in the preliminary elements it was impossible to decide on which branch of the curve they should lie. For this reason, and also because there is generally a systematic difference between radial velocities obtained at two observatories, the period was determined from the Ottawa observations alone. As it turned out, the inclusion of the Yerkes Observatory plates would have yielded practically the same result, for the final orbit gives a mean residual for their plates of +3.1 kilometres, which is just about the difference usually found between measures made by the two observatories. Seventeen normal places were formed as given. It is unfortunate that the negative minimum is not more strongly determined, but the lack of observations is due

to the star and sun being in conjunction at that time. If the star were followed for another revolution this defect could be remedied. In order to determine the period accurately, the star should be observed at some epoch in the future which can be so chosen as to define the negative minimum as well.

The preliminary elements chosen follow the normal places and the successive steps in the least-square solution are recorded.

NORMAL PLACES

No.	Julian Day	Phase from Preliminary T	Velocity	Weight	O-C Preliminary	O-C Final
1	2,420,077.8	408.8	+19.3	.3	+1.2	+1.4
2	429.4	160.4	-16.4	.3	-3.8	-3.4
3	485.2	216.2	-2.3	.4	+0.6	+1.4
4	443.8	111.8	+19.2	.8	+0.9	+1.3
5	467.9	198.9	+14.0	.6	-2.3	-1.6
6	492.1	523.1	+12.5	.8	-1.1	-0.5
7	531.4	565.4	+8.3	.4	+1.6	+2.4
8	571.6	602.6	-4.3	.4	-1.6	-1.3
9	609.0	640.0	-12.4	.2	+1.3	-0.6
10	730.3	101.3	-21.6	.4	-0.1	-0.9
11	761.4	132.4	-16.2	.4	+0.8	+0.4
12	800.4	171.4	-9.2	.6	+1.2	+1.0
13	839.9	210.9	-3.5	.4	+0.3	+0.2
14	879.4	250.4	+2.1	.5	-0.2	-0.2
15	920.7	291.7	+6.9	.9	-0.9	-1.0
16	948.4	319.4	+11.1	.8	+0.5	+0.4
17	987.2	358.2	+11.9	.2	+0.2	+0.3

PRELIMINARY ELEMENTS

$$\begin{aligned}
 P &= 660 \text{ days} \\
 T &= \text{J. D. } 2,420,629 \\
 \omega &= 135^\circ \\
 e &= 0.20 \\
 K &= 21.5 \\
 \gamma &= -0.16 \\
 \mu &= 0^\circ.545455
 \end{aligned}$$

OBSERVATION EQUATIONS

	x	y	z	p	q	r	$-n$	Weight
1	1.000	- .916	+ .531	+ .152	- .629	- .347	- 1.21	0.3
2	1.000	- .565	+ 1.007	+ .765	- .787	- .391	+ 3.78	0.3
3	1.000	- .428	+ .639	+ .859	- .734	- .326	- 0.62	0.4
4	1.000	+ .859	- .711	- .135	- .001	- .001	- 0.90	0.8
5	1.000	+ .769	+ .077	- .557	+ .360	- .058	+ 2.34	0.6
6	1.000	+ .651	+ .528	- .747	- .571	+ .079	+ 1.39	0.8
7	1.000	+ .319	+ 1.105	- 1.029	+ 1.000	+ .095	- 1.61	0.4
8	1.000	- .120	+ .901	- 1.141	+ 1.295	+ .074	+ 1.55	0.4
9	1.000	- .616	- .133	- 1.021	+ 1.250	+ .025	- 1.31	0.2
10	1.000	- .902	+ .381	+ .381	- .571	+ .058	+ 0.10	0.4
11	1.000	- .783	+ .809	+ .625	- .738	+ .097	- 0.80	0.4
12	1.000	- .475	+ .989	+ .801	- .787	+ .135	- 1.18	0.6
13	1.000	- .167	+ .701	+ .859	- .741	+ .157	- 0.26	0.4
14	1.000	+ .114	+ .247	+ .826	- .661	+ .165	+ 0.18	0.5
15	1.000	+ .370	- .308	+ .719	- .559	+ .163	+ 0.89	0.9
16	1.000	+ .514	- .606	+ .614	- .485	+ .155	- 0.52	0.8
17	1.000	+ .693	- .896	+ .410	- .355	+ .127	- 0.17	0.2

Where $x = d\gamma$

$y = dK$

$z = Kde$

$p = Kd\omega$

$q = \frac{K\mu}{(1-e^2)^{\frac{3}{2}}} dT$

$r = \frac{1000 K}{(1-e^2)^{\frac{3}{2}}} d\mu$

NORMAL EQUATIONS

$$\begin{aligned}
 +8.400r &+ 1.001y &+ 1.931z &+ 1.459p &- 1.465q &+ 0.402r &+ 1.382 = 0 \\
 +3.003y &- 1.638z &- 1.035p &+ 0.986q &+ 0.297r &+ 1.433 = 0 \\
 &+ 3.708z &- 0.070p &- 0.026q &- 0.111r &+ 0.903 = 0 \\
 &&&+ 4.367p &- 4.050q &+ 0.017r &- 1.267 = 0 \\
 &&&&&+ 3.890q &+ 0.028r &+ 1.052 = 0 \\
 &&&&&&+ 0.223r &- 0.153 = 0
 \end{aligned}$$

Whence $x = +0.013$ or $d\gamma = +0.01$ km.
 $y = -0.969$ $dK = -0.97$ km.
 $z = -0.625$ $de = -0.029$
 $p = +0.197$ $d\omega = +0^\circ.52$
 $q = +0.169$ $dT = +0.78$ day
 $r = +1.606$ $d\mu = +0^\circ.004025$

FINAL ELEMENTS

$$\begin{aligned}
 P &= 655.16 \pm 5.26 \text{ days} \\
 T &= J. D. 2,420,629.78 \pm 0.56 \text{ days} \\
 \omega &= 135^\circ.52 \pm 5.2^\circ \\
 e &= 0.171 \pm 0.026 \\
 K &= 20.53 \text{ km.} \pm 0.57 \text{ km.} \\
 \gamma &= -0.15 \text{ km.} \pm 0.35 \text{ km.} \\
 a \sin i &= 182,300,000 \text{ km.} \\
 \frac{m_1^3 \sin^3 i}{(m + m_1)^2} &= 0.56 \odot
 \end{aligned}$$

THIRD BODY

It will be noticed that the binary character was announced from three plates, which the orbit shows to fall near the epoch of positive maximum (cf. radial velocity curve)*. In the interval covered by these plates the variation should be small and the velocities increasing. One is forced to conclude that there is another period superimposed upon the one found, or else that the announced variation means nothing. The spectrograms have been remeasured at the Yerkes Observatory, and Professor Frost finds no reason to doubt that the range indicated by the velocities is real.

We have been unable to find a period for this variation. Several periods would harmonize some of the larger residuals and so reduce the sum Σp^2 a little, but the fact that there are several which look equally good (or bad) is enough to render one suspicious of the reality of any. No period found would satisfy the residuals from the main orbit well enough to warrant its adoption.

Considering the elements of the system as published, the probable error of a single observation is 3.5 kilometres. This is larger than we usually find for good line stars by about 20 per cent.

In the *Astrophysical Journal*, March, 1915, Dr. Schlesinger published a method for determining the periodicity in a series of observations from a consideration of the distribution of the observed velocities. Thus, for example, if we are dealing with a circular orbit, more velocities should be found near positive maximum and negative minimum than near the γ line. If the residuals are due entirely to error of observation, the velocities around the γ line will be most numerous. Curves are published giving the expected distribution of the velocities for orbits of various forms. In these curves some allowance has been made for the effect of probable error of measurement upon the results.

Some time previous to the publication of this article, the writer tried to make use of the same method in getting a period for the binary 12 Lacertae. In this case the velocities were distributed very much as one would expect if the observed range were due to error of measurement alone and yet, since the total observed range was over sixty kilometres, there could be no doubt of the binary nature of the star. The conclusion

*The normal place from the observations taken at the Yerkes Observatory is indicated by a circle and cross.

was that the expected distribution was affected very much by errors of measurement. In applying the method to the present star I have taken some care to find out just what distribution of velocities we should expect when both factors are taken into account.

Let us suppose that we are dealing with a circular orbit whose range is from plus ten kilometres to minus ten kilometres, and that the probable error of measurement is r . Let the velocities, if there were zero error of measurement, which would lie between the limits plus ten to plus nine be denoted by a , plus nine to plus eight be denoted by b , plus eight to plus six by c , , plus two to zero by f . Let those from zero to minus two be denoted by f' , and so on till those between minus nine and minus ten are denoted by a' . Secondly, let us suppose the range in the observations is due to error of measurement alone, and let the velocities which would lie between the limits zero to plus two be denoted by I , between the limits plus two to plus four by I_2 , etc. to infinity. To distinguish the negative from the positive add the subscript minus. Thus I_- indicates the velocities lying between minus two and zero.

The numbers a , b , c , are easily computed analytically or graphically. The graphical method has the advantage that it can be applied to any radial velocity curve, while the analytic expression for a , b , c , for other than circular orbits is very cumbersome. The values of I , I_2 , I_3 , can be readily computed for any assumed probable error.

Now let all the velocities a , b , c , etc., be operated on by errors of measurement, which errors are distributed as indicated by I , I_2 , I_3 , etc. The product of a and I (which may be written a_I) will yield velocities lying between +9 and +12, and if we have chosen our interval small enough we may regard the velocities as distributed uniformly within this interval, so that $\frac{2}{3} a_I$ will be the number lying between +12 and +10. $\frac{1}{3} a_I$ will also lie between the same limits, and if we select all these combinations we will obtain the following series as giving the number of velocities between +12 and +10.

$$\begin{aligned} & \frac{2}{3} a_1 + \frac{1}{3} a_2 + \frac{1}{3} b_1 + \frac{2}{3} b_2 + \frac{1}{2} c_2 + \frac{1}{2} c_3 \\ & + \frac{1}{2} d_3 + \frac{1}{2} d_4 + \frac{1}{2} e_4 + \frac{1}{2} e_5 + \frac{1}{2} f_5 + \frac{1}{2} f_6 \\ & + \frac{1}{2} f'_6 + \frac{1}{2} f'_7 + \frac{1}{2} c'_7 + \frac{1}{2} c'_8 + \frac{1}{2} d'_8 + \frac{1}{2} d'_9 \\ & + \frac{1}{2} c'_9 + \frac{1}{2} c'_0 + \frac{2}{3} b'_0 + \frac{1}{3} b'_1 + \frac{2}{3} a'_1 + \frac{1}{3} a'_0 \end{aligned}$$

To obtain the velocities lying between +10 to +8 all that it is necessary to do is to depress the subscripts by one, $\frac{2}{3} a_1$ becomes $\frac{2}{3} a_{1-}$, $\frac{1}{3} a_2$ becomes $\frac{1}{3} a_{2-}$, etc. To obtain the velocities lying between +14 and +12 increase the subscripts by one. The number of velocities lying between limits x and $x+2$ are obtained by increasing the subscripts $\underline{x-10}$.

2

These series have been evaluated for $r = 2, 4, 6.66, 10$ kilometres, i.e., for the ratios of probable error of measurement to total range equal $\frac{1}{10}, \frac{1}{5}, \frac{1}{3}, \frac{1}{2}$. The results are tabulated below and represented graphically in fig. 1.

Velocities in km.	$r=2$ km.	$r=4$ km.	$r=6\cdot6$ km.	$r=10$ km.
From + 0 to + 2	.0680	.0753	.0637	.0479
" + 2 to + 4	.0795	.0748	.0623	.0474
" + 4 to + 6	.0833	.0723	.0591	.0461
" + 6 to + 8	.0878	.0671	.0554	.0441
" + 8 to + 10	.0793	.0604	.0503	.0415
" + 10 to + 12	.0571	.0495	.0442	.0387
" + 12 to + 14	.0295	.0380	.0384	.0356
" + 14 to + 16	.0104	.0266	.0318	.0321
" + 16 to + 18	.0025	.0173	.0258	.0284
" + 18 to + 20	.0004	.0099	.0201	.0249
" + 20 to + 22		.0053	.0150	.0215
" + 22 to + 24		.0024	.0114	.0185
" + 24 to + 26		.0009	.0079	.0153
" + 26 to + 28		.0002	.0052	.0124
" + 28 to + 30		.0001	.0031	.0100
" + 30 to + 32			.0019	.0079
" + 32 to + 34			.0010	.0061
Sum	.4978	.5004	.4963	.4784

In the tabulated results the numbers are given to four decimal places, and as each was obtained by the summation of several factors, each of which was only computed to four places, the last figure is not in general correct. The sum of the columns given at the bottom if extended to infinity should total one-half, so that the addition of the columns serves as a partial check on the work.

In the curves the abscissae are kilonettes, any point plus x indicates velocities lying between plus $x-1$ and plus $x+1$ and the corresponding ordinate shows what fraction of all the velocities will lie between these limits. It will be noticed that curves for the ratios $\frac{1}{2}$, $\frac{1}{3}$, $\frac{1}{5}$ are very similar to error curves.




Fig. 1

Suppose now we are working with such a binary as the one indicated, namely, a range plus ten to minus ten, and error of measurement r kilometres. Let us suppose, further, that we are unable to get a period, and treat the residuals as errors, how much in excess of the true value of r would the value we obtain be? The results we have tabulated for the distribution of the velocities enable us to compute this with a fair degree of accuracy, and we obtain that, if r equal 2 kilometres the computed r will be 5.1, if r equal 4 kilometres the computed r will be 6.1, and if r equal 6.6 kilometres the computed r will be 8.2. The computation becomes a little uncertain for r equal 10 on account of some velocities lying beyond the computed values.

To apply the results to χ Aurigæ, we have computed the distribution of the velocities on two bases. In the first the residuals are supposed to be due to error of measurement alone, in the second they are supposed to be due to a probable error of measurement 2.84 kilometres, combined with a binary whose eccentricity is zero and range 8.52 kilometres. In the table below, the first line gives the distribution of the observed residuals, the second, the computed distribution on hypothesis one, and the third the computed distribution on hypothesis two.

	km. 0-1	km. 1-2	km. 2-3	km. 3-4	km. 4-5	km. 5-6	km. 6-8	km. 8-10	km. 10-∞
1	14	15	11	14	6	4	11	7	6
2	13	13	12	11	9	8	11	5	6
3	13	13	12	11	9	8	11	6	5

The two hypotheses give almost identical distributions when the number of observed velocities is limited, in this case 88. In other words, an inspection of the manner in which the velocities are distributed would not enable us to detect the presence of a velocity variation three times the probable error of measurement.

The above results are deduced for circular orbits only. They have to be modified slightly for orbits of high eccentricity, but in these also the effect of error of measurement is to mask the presence of the orbital variation.

The only evidence that we have of a third body in the system of χ Aurigae is the magnitude of the probable error obtained for a single velocity. If we assume that it should be 2.8, then it would be necessary to assume also a range of 8.5 kilometres in order to obtain a distribution of residuals to agree with the observed.

CALCIUM LINES

In reducing the measures of the different plates the two calcium lines were not included with the others, for evidence was found showing they would yield a smaller range for the curve. After the completion of the orbit as given in the preceding pages, the velocities from the H and K lines were grouped into ten normal places as follows.

NORMAL PLACES FOR H AND K LINES

	Julian Day	Phase from Final T	Velocity	Weight	O-G
1.....	2,420,104.4	129.8	- 9.5	1.0	-2.3
2.....	151.3	176.7	- 0.8	0.8	+2.5
3.....	202.4	227.8	+ 0.5	0.9	-0.4
4.....	249.9	275.3	+ 7.6	1.0	+3.2
5.....	277.0	302.5	+ 5.2	1.3	-0.9
6.....	302.9	328.3	+ 10.3	1.2	+2.8
7.....	411.8	437.2	+ 9.9	1.3	-0.6
8.....	473.4	498.8	+ 7.2	1.4	-1.9
9.....	502.1	527.5	+ 7.2	1.4	-0.1
10.....	571.2	596.6	+ 0.6	1.5	+0.7

Preliminary elements to satisfy these places were selected, the period, eccentricity and ω were taken as previously found for the other lines. A solution was carried through for γ and K with the result,

$$\begin{aligned}\gamma &= + 1.5 \text{ km.} \pm 0.46 \\ K &= 10.47 \text{ km.} \pm 0.85\end{aligned}$$

The radial velocity curve for the H and K lines is given on the same diagram as that for the other lines and is drawn with the finer line. The difference between the

velocity of the system obtained from the *H* and *K* lines on the one hand, and the rest of the lines on the other, would disappear if the assumed values of wave-lengths for the calcium lines were increased 0.022 tenth-metres.

The behaviour of the calcium lines in stellar spectra has been a subject for much speculation and investigation since the orbit of δ Orionis was published by Hartmann.¹ The chief facts so far brought out may be condensed as follows:—

1. In eight stars the calcium lines are constant, or nearly so, and are sharp and narrow, while the other lines are more or less diffuse and show a large range in velocity. These stars are given below.

Name	R.A. 1900	Type	Range of Broad Lines	Remarks
	h m		km.	
σ Persei.....	3 38.0	B1	224	Lick Obs. Bull. 181
η Orionis.....	5 19.4	B1	290	" "
ψ Orionis.....	5 21.6	B2	288	" "
δ Orionis.....	5 26.9	B	216	" "
VV Orionis.....	5 28.5	B2	264	P.A.O., Vol. III, No. 21
θ_1 Orionis.....	5 30.5	B1	220	Lick Obs. Bull. 181
β Scorpii.....	15 59.6	B1	251	P.A.O., Vol. II, No. 14
Boss 6142.....	23 51.0	Oep	230	D.O. (unpublished)

All the stars in this list are of type B2 or earlier, and, moreover, if we form a list of all such stars whose orbits have been determined and the *K* line measured, we find only one, ν Orionis, showing a large range and not included above. Future orbits may add other exceptions.

2. In several stars, the calcium lines are known to vary differently from the other lines and in four cases, orbits have been determined from the *H* and *K* lines, *i.e.*, for ξ Persei², type Oe5; 9 Camelopardalis³, type B; χ Aurigæ, type B1; 12 Lacertæ⁴, type Oe5. In the last two the amplitude is probably about one-half as large for the calcium lines as for the others. There can be little doubt but that future investigation will reveal all gradations from those in which the *H* and *K* lines are constant to those where they have the same oscillation as the other lines.

3. In three stars the calcium lines are sharp and narrow and are shifted about one-half an Ångström unit to the violet. These stars are,— ξ Persei, type Oe5; ϕ_1 Orionis⁵, type B; ρ Leonis⁶, type B. There is evidence in these cases that it is the hydrogen

¹Ap. J., Vol. XIX, p. 268, 1904.

²Dom. Obs. Publ., Vol. I, p. 355.

³Ap. J., Vol. XXXVII, p. 1, 1913.

⁴R.A.S.C.Jr., November, 1915.

⁵Ap. J., Vol. XXX, p. 63, 1909.

⁶Dom. Obs. Publ., Vol. I, p. 337.

and helium lines that are shifted to the red and that the calcium lines may be in their normal positions.

In ξ Persei, the hydrogen and helium lines are broad and no orbit has been determined from them, although they seem to show a large range. The mean velocity is about plus sixty-five kilometres. An orbit has been determined from the H and K lines, and the velocity of the system is plus fifteen kilometres. The component of the solar motion away from this star is about plus six kilometres, so that unless the star has an abnormally high velocity for an early B-type star we must regard the K lines as giving the true velocity of the star.

In the case of ϕ_1 Orionis, no orbit has been published, but if we take the mean of the velocities as representing final values, the calcium lines give plus nineteen, the hydrogen and helium lines plus forty, as compared with a computed velocity from the solar motion plus fifteen. Here again it is the velocity from the H and K lines that seems the most probable.

No orbit has been published for ρ Leonis, but the mean velocities are, for the calcium lines plus ten, for the other lines plus forty-three, for the solar motion plus five.

4. In several stars (κ Cassiopeia⁵, type B; 9 Camelopardalis, type B; ϵ Orionis⁶, type B), the calcium lines are sharp and the other lines diffuse, but there does not seem to be much difference between the mean velocities as derived from the two sets of lines. The difference between these stars and those in 2 is only one of degree.

5. In many stars of early B-type, all the lines are very diffuse and in some cases the spectrum is almost continuous, so that little can be said concerning the behaviour of the calcium. α Virginis and δ Scorpii are representative.

6. There seems to be no exception to the rule, that when the calcium lines are sharp and narrow and the other lines broad, the star exhibits a variable radial velocity.

Three theories have been advanced to explain the observations. In the first, the calcium lines are supposed to be due to masses of cooler gas between the observer and the binary. In the second, the gas is supposed to surround the star or stars, and in the third, the phenomena are ascribed to anomalous dispersion.

The first hypothesis accounts for the sharp and narrow character of the H and K lines in class 1 and the fact that they yield constant velocity. Why should the calcium clouds always lie in front of a star of type B2 or earlier? We have examined several later B-type stars without finding anywhere H and K are stationary. There may be some stars later than B2 discovered which will show the effect, but the phenomenon seems to be one connected with early spectral type. In all cases in class 1 the velocities yielded by the calcium lines are very near the velocity of the system. This fact, however, does not argue strongly either against or for the first hypothesis on account of the very low velocities of the early B-type stars. The stars given under 3 require an explanation for the hydrogen and helium lines rather than for the calcium. The phenomenon is too complex to be accounted for by an hypothesis which allows so little latitude for adjustment.

⁵Unpublished results (Harper).
⁶Ap. J., Vol. XXIX, p. 236, 1908.

The second hypothesis may be presented as follows:—



Fig. 2

Let A and B (fig. 2) be two stars or the two nuclei of a binary, and let the gases be condensed about these. It is known that calcium occurs at very high levels in the sun, and let us suppose that the equidensity layers in the calcium vapour about the two stars are nearly spheres. The calcium cloud will rotate about the common centre of gravity of the two stars in the same period as they revolve about each other. If A is the primary star and only one set of lines is observed on the plate, we need concern ourselves only with the light from A. The light from the photosphere gives the continuous spectrum, and as it passes through the cooler gases, hydrogen, helium and calcium, has lines from these elements absorbed. The absorption of the calcium takes place where the element is very rare, which accounts for the narrow character exhibited by these lines. The absorption of helium and hydrogen must be supposed to take place at a much lower level. Assume that it is permissible to speak of effective levels at which this absorption occurs, at a height D for the hydrogen and helium and at a height F for the calcium. At the time when the primary is moving toward us with the greatest velocity, the points at which the calcium absorption is taking place are rotating then as always about the point C and so are moving nearly across the line of sight. Moreover, the different points are all moving at about the same rate toward the observer, thus maintaining the narrow character of the lines. The points at which the hydrogen and helium absorption occurs, exhibit a wide range in velocity and give the wide character of the lines and the high range if the inclination is nearly zero. If the inclination is nearer to ninety degrees, the hydrogen and helium lines would be sharper and give a smaller range. This fits the case of 12 Lacertæ and X Aurigæ. There are doubtless other factors not understood which may contribute to the width of the lines observed in other cases. If the calcium cloud were extensive enough the calcium lines would remain nearly stationary, at least the amplitude would be small enough to be masked by errors of measurement. As the cloud condensed, the amplitude observed in the H and K lines would increase, the exact range depending on the degree of condensation of the calcium.

This theory is capable of considerable adjustment⁹ to meet varying demands, and suggests future types of variations that should be discovered.


The third hypothesis ascribes the wide character of the lines in these stars to anomalous dispersion effects. The calcium lines are sharp and narrow, due to the small traces of these elements existing in the star's atmosphere. It is not intended to pass on the merits or demerits of this theory, but the fact that it does not satisfactorily account for the periodicity in the radial velocities has prevented its receiving wide acceptance.

⁹Ap. J. Vol. XXXVII, p. 1, 1913.

SUMMARY

1. An orbit has been determined from 88 measures of radial velocity of α Aurige.
2. The amplitude shown by the calcium lines is about one-half that for the other lines of the spectrum.
3. An analysis of the different cases so far known where the calcium lines behave differently from the other lines shows that the phenomenon is one of type, and is best explained by the presence of calcium in a cloud surrounding the binary. The absorption of calcium takes place at a much higher level than that of the other elements.
4. The residuals given from the simple elliptic orbit were examined to ascertain if there might be a third body present. No secondary period was found, but the star should be observed with higher dispersion to test this point further.
5. The method of detecting periodicity in a set of observations from a consideration of the distribution of the velocities was tested. The value of this method is greatly interfered with by the effects of errors of measurement. If the ratio of error of measurement to range in the binary is as great as one-fifth, the resulting distribution of velocities is very much the same as would result from errors alone. If the ratio is as large as one-third, the distribution is almost identical. It is in these very cases that the method is most likely to be needed.

Dominion Observatory
Ottawa
June, 1916.



NOTE ON THE SPECTROSCOPIC BINARY 12 LACERTÆ
BY REYNOLD K. YOUNG, PH.D.

The elements of this binary were published in the *Publications of the Dominion Observatory*, Vol. II, No. 3. An examination of the velocities given, shows the *K* line to have a smaller range than the other lines in the spectrum. If the calcium lines alone had been used, the amplitude of the curve would have been about one-half that deduced from the lines for the other elements.

In the following table the differences between the velocities as given by the *K* line and those given by the other lines are grouped according to phase. The period is divided into four parts. The first column includes the positive maximum and the third the negative minimum. The second and fourth columns include the zero axis of the system.

Plate No.	Difference in km.						
5670	-12.0	5642	+ 1.4	5623	+ 7.3	5630	+ 1.4
5735	-22.5	5720	+ 5.5	5692	+ 7.5	5680	+ 3.0
6244	- 3.0	5800	+ 6.9	5728	+ 5.7	5711	+ 1.3
6276	- 0.9	6198	+12.1	5792	+ 8.9	5787	- 4.1
6328	- 3.9	6335	+13.7	6270	+ 7.9	6262	+17.1
6369	+ 5.3	6349	+16.2	6281	+ 1.8	6291	+19.4
6409	- 9.9	6501	- 2.0	6342	+ 3.6	6308	- 2.6
6426	- 4.2	6992	- 2.0	6531	+ 6.1	6356	+11.0
6497	- 2.4	6998	+ 3.8	6542	- 4.1	6454	- 5.4
6521	+12.6	7033	- 9.4	7001	+14.8	7023	- 8.1
6557	+ 0.4	7043	+11.9	7020	+14.2	7178	+ 0.3
7026	- 1.9	7183	+ 9.9	7177	+22.5	7210	+16.0
7176	-13.3	7220	- 4.2	7214	+16.9	7244	- 4.4
7182	+ 1.2			7243	+ 5.2		
7201	-12.2			7251	+ 9.9		
7211	- 9.8			7266	+ 6.7		
7249	- 2.3			7270	+15.0		
7268	+ 4.1			7293	- 2.2		
7302	- 2.5			7300	+10.0		
7310	-16.8			7313	+28.4		
7314	-11.0						
Mean	- 5.2	Mean	+ 4.9	Mean	+ 9.7	Mean	+ 3.5

The weighted mean of all the residuals is positive, due to the value of the wavelength of *K* used. If we remove this effect, the resulting means for columns 1, 2, 3, 4 would read -7.3 km., -0.2 km., +7.6 km., +1.4 km. The semi-amplitude of the velocity curve as found for all the lines was 16.92 km. The amplitude for the *K* line would be about 7.5 km. less.

Dominion Observatory

Ottawa

May, 1916.



3 9334 00061480 8

CAL
MT 403
P04-01

15 JUN 1999 A2B

130007

